

CLINICAL TRIALS AND OBSERVATIONS

Novel use Of Hydroxyurea in an African Region with Malaria (NOHARM): a trial for children with sickle cell anemia

Robert O. Opoka,¹ Christopher M. Ndugwa,¹ Teresa S. Latham,² Adam Lane,² Heather A. Hume,³ Phillip Kasirye,¹ James S. Hodges,⁴ Russell E. Ware,^{2,*} and Chandy C. John^{5,*}

¹Department of Paediatrics and Child Health, Makerere University, Kampala, Uganda; ²Division of Hematology, Department of Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, OH; ³Département de Pédiatrie, Université de Montréal Service d'Hématologie/Oncologie, Montréal, QC, Canada; ⁴Division of Biostatistics, School of Public Health, University of Minnesota, Minneapolis, MN; and ⁵Ryan White Center for Infectious Diseases and Global Health, Department of Pediatrics, University of Indiana, Indianapolis, IN

Key Points

- Compared with placebo, hydroxyurea did not increase the incidence or severity of malaria events in Ugandan children with SCA.
- Hydroxyurea provided significant clinical and laboratory benefits, suggesting it will be safe and effective across sub-Saharan Africa.

Hydroxyurea treatment is recommended for children with sickle cell anemia (SCA) living in high-resource malaria-free regions, but its safety and efficacy in malaria-endemic sub-Saharan Africa, where the greatest sickle-cell burden exists, remain unknown. In vitro studies suggest hydroxyurea could increase malaria severity, and hydroxyurea-associated neutropenia could worsen infections. NOHARM (Novel use Of Hydroxyurea in an African Region with Malaria) was a randomized, double-blinded, placebo-controlled trial conducted in malaria-endemic Uganda, comparing hydroxyurea to placebo at 20 ± 2.5 mg/kg per day for 12 months. The primary outcome was incidence of clinical malaria. Secondary outcomes included SCA-related adverse events (AEs), clinical and laboratory effects, and hematological toxicities. Children received either hydroxyurea (N = 104) or placebo (N = 103). Malaria incidence did not differ between children on hydroxyurea (0.05 episodes per child per year; 95% confidence interval [0.02, 0.13]) vs placebo (0.07 episodes per child per year [0.03, 0.16]); the hydroxyurea/placebo malaria incidence rate ratio was 0.7 ([0.2, 2.7]; $P = .61$). Time to infection also did not differ significantly between treatment arms. A composite SCA-related clinical outcome (vaso-occlusive painful crisis, dactylitis, acute chest syndrome, splenic sequestration, or blood transfusion) was less frequent with hydroxyurea (45%) than placebo (69%; $P = .001$). Children receiving hydroxyurea had significantly increased hemoglobin concentration and fetal hemoglobin, with decreased leukocytes and reticulocytes. Serious AEs, sepsis episodes, and dose-limiting toxicities were similar between treatment arms. Three deaths occurred (2 hydroxyurea, 1 placebo, and none from malaria). Hydroxyurea treatment appears safe for children with SCA living in malaria-endemic sub-Saharan Africa, without increased severe malaria, infections, or AEs. Hydroxyurea provides SCA-related laboratory and clinical efficacy, but optimal dosing and monitoring regimens for Africa remain undefined. This trial was registered at www.clinicaltrials.gov as #NCT01976416. (*Blood*. 2017;130(24):2585-2593)

dactylitis, acute chest syndrome, splenic sequestration, or blood transfusion) was less frequent with hydroxyurea (45%) than placebo (69%; $P = .001$). Children receiving hydroxyurea had significantly increased hemoglobin concentration and fetal hemoglobin, with decreased leukocytes and reticulocytes. Serious AEs, sepsis episodes, and dose-limiting toxicities were similar between treatment arms. Three deaths occurred (2 hydroxyurea, 1 placebo, and none from malaria). Hydroxyurea treatment appears safe for children with SCA living in malaria-endemic sub-Saharan Africa, without increased severe malaria, infections, or AEs. Hydroxyurea provides SCA-related laboratory and clinical efficacy, but optimal dosing and monitoring regimens for Africa remain undefined. This trial was registered at www.clinicaltrials.gov as #NCT01976416. (*Blood*. 2017;130(24):2585-2593)

Introduction

Sickle cell anemia (SCA) is a life-threatening hematological disorder and among the world's most prevalent hereditary diseases, with more than 300 000 affected babies born each year.¹ The vast majority of these births occur in sub-Saharan Africa, where an estimated 50% to 90% of children with SCA will die by age 5 years, often without an established diagnosis.² The heterozygous sickle gene mutation confers a strong survival advantage against malaria,³ which explains why the allele frequency is highest in malaria-endemic regions of Africa.¹

Hydroxyurea has proven laboratory and clinical efficacy for both children and adults with SCA.⁴⁻⁷ Its mechanisms of action are multiple and incompletely understood, but fetal hemoglobin (HbF) induction in erythroid cells is critical for the inhibition of intracellular sickling. Treatment also has salutary effects on blood cell adhesion, morphology, and rheology.⁸ Further, hydroxyurea is a safe drug for SCA, with low

incidence of treatment-related toxicity and no serious long-term effects observed to date. Hydroxyurea is currently approved by the US Food and Drug Administration for adults with severe symptoms and by the European Medicines Agency for affected adults and children above age 2 years. Based on a large and compelling body of evidence accumulated over the past 30 years, evidence-based guidelines published by the National Heart, Lung and Blood Institute of the National Institutes of Health strongly recommend wider usage, including offering treatment to infants as young as 9 months of age.⁹

The benefits of hydroxyurea treatment of children with SCA would be greatest in countries within sub-Saharan Africa, if hydroxyurea were found to be safe and have its predicted efficacy in these high-burden areas. However, the effects of hydroxyurea on the presentation and clinical course of malaria must also be considered. Hydroxyurea

Submitted 3 June 2017; accepted 18 August 2017. Prepublished online as *Blood* First Edition paper, 19 October 2017; DOI 10.1182/blood-2017-06-788935.

*R.E.W. and C.C.J. are joint senior authors.

There is an Inside *Blood* Commentary on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

© 2017 by The American Society of Hematology

Table 1. NOHARM AE definitions

	Definition
Clinical AE terminology	
Vaso-occlusive pain crisis/dactylitis	Vaso-occlusive pain crisis: acute pain and tenderness in an area of the body, with or without swelling, with no other diagnostic explanation. Dactylitis: vaso-occlusive crisis (acute pain, tenderness, and swelling) localized to hands or feet.
Pneumonia/acute chest syndrome	Pneumonia: history of fever or measured axillary temperature $\geq 37.5^{\circ}\text{C}$, with tachypnea and cough. Acute chest syndrome: signs of pneumonia plus chest pain and/or tenderness.
Clinical sepsis	Measured fever and ill appearance, requiring IV antibiotics.
Acute splenic sequestration	Increase in splenic size from last physical examination, accompanied by a decrease in hemoglobin of ≥ 2 g/dL.
Upper respiratory infection	Child with general well appearance with rhinorrhea, nasal congestion, or cough.
Gastrointestinal related	Diarrhea, vomiting, constipated, intestinal obstruction.
Malaria	Measured fever (axillary temperature $\geq 37.5^{\circ}\text{C}$) or fever by history and <i>Plasmodium</i> species infection on blood smear.
Other infection	Other infections, diagnosed clinically.
Other (eg, injury)	Other diseases not included above diagnosed during visits for illness.
Laboratory AEs*	
Anemia	Hemoglobin < 6 g/dL
Reticulocytopenia	ARC $< 80 \times 10^9/\text{L}$ and hemoglobin < 7 g/dL
Neutropenia	Absolute neutrophil count (ANC) $< 1.0 \times 10^9/\text{L}$
Thrombocytopenia	Platelet count $< 80 \times 10^9/\text{L}$
Elevated AST/ALT	AST > 150 IU/L, ALT > 150 IU/L
Elevated bilirubin	Total bilirubin > 5 mg/dL

The diagnosis of sickle-related clinical events followed published definitions,²⁴ with modifications such as pneumonia and clinical sepsis. ALT, alanine transferase; AST, aspartate transferase.

*Laboratory AE definitions represent the values necessary for a grade 2 event.

may directly affect several factors related to the pathogenesis of severe malaria, with potentially deleterious consequences. Some in vitro and animal studies suggest that hydroxyurea increases endothelial intracellular adhesion molecule-1 (ICAM-1) expression,¹⁰ which could enhance parasite adhesion to endothelium,¹¹ and also increases tumor necrosis factor α levels,¹² both of which are associated with increased malaria severity and death.¹³ Other studies have challenged these findings,^{14,15} but definitive human data are lacking. If hydroxyurea induces effects that shift the clinical course in SCA from uncomplicated to severe malaria, this could increase mortality despite benefits for the underlying SCA.¹⁶ In addition, in low-resource settings, hydroxyurea may cause neutropenia that could lead to increased severity of the many bacterial infections commonly observed in African children with SCA.¹⁷

In contrast to these adverse effects, it is possible that hydroxyurea could have beneficial effects against malaria because HbF, which is increased by hydroxyurea, inhibits *Plasmodium falciparum* growth in vitro.¹⁸ Hydroxyurea treatment also generates nitric oxide,¹⁹ which protects against severe malaria in animals^{20,21} and humans.²² With these contrasting potential mechanisms, the risks and benefits of hydroxyurea in a malaria endemic setting remain unknown, and many critical questions remain regarding the safety, feasibility, and efficacy of hydroxyurea for children with SCA living in malaria-endemic settings within sub-Saharan Africa.

For these reasons, we conducted a prospective randomized double-blinded placebo-controlled clinical trial, the Novel use Of Hydroxyurea in an African Region with Malaria (NOHARM), in young Ugandan children with SCA, to determine the safety and efficacy of hydroxyurea in a malaria-endemic region. NOHARM is registered at www.clinicaltrials.gov as #NCT01976416.

Methods

Study design

A detailed description of the study site, design, and procedures was previously published.²³ Briefly, NOHARM was a randomized double-blinded placebo-controlled clinical trial. Because of concerns about the risks of severe

malaria and infection with hydroxyurea treatment, the consensus among local experts and the local institutional review board supported a placebo-controlled trial, despite the expected SCA-related treatment benefits in children with SCA. It was agreed, however, that study participants should be given the opportunity to receive subsequent open-label hydroxyurea, if no danger signal was observed from the blinded study treatment. NOHARM was conducted at Mulago Hospital Sickle Cell Clinic in Kampala, Uganda. Kampala has seasonal malaria transmission, with high outpatient and inpatient burdens. Protocol approval was obtained from the institutional review boards of the Makerere University School of Medicine, Indiana University, University of Minnesota, and Cincinnati Children's Hospital, as well as the Uganda National Drug Authority and Uganda National Council for Science and Technology.

Study participants

Children receiving care at Mulago Hospital Sickle Cell Clinic were eligible if they were 1.00 to 3.99 years of age and living within 50 km of the clinic. Study inclusion criteria included confirmed SCA, weight ≥ 5.0 kg, and willingness to comply with study procedures. Children with severe malnutrition or known chronic medical conditions, current hydroxyurea treatment, or blood transfusion in the previous 30 days were excluded. Parents of the study participants gave written informed consent.

Randomization and masking

Children meeting inclusion criteria completed enrollment and after screening, were randomized 1:1 to hydroxyurea or placebo by the Data Coordinating Center at Cincinnati Children's Hospital as described.²³ Study allocations were randomly assigned by the statistician, using computer-generated sequences in block files of 8 participants. Study treatments were supplied with A or B labels, and the clinical coordinating center staff, study pharmacists, families and caregivers, and all but 2 members of the Data Coordinating Center were masked to treatment allocation.

Medications

Study treatment (Addmedica, Paris, France) included oral hydroxyurea (Siklos) as 1000-mg scored tablets and 100-mg dispersible tablets, or placebo tablets of identical size and appearance. Treatment was administered once daily at 20 ± 2.5 mg/kg for 12 months, with dose adjustments in both arms for weight gain and hematological toxicities.²³




Figure 1. CONSORT diagram for the NOHARM trial.

Procedures

All participants received standard care for SCA including folic acid, penicillin prophylaxis, and pneumococcal vaccination. For malaria prophylaxis, children received insecticide-treated mosquito nets and monthly sulphadoxine-pyrimethamine. Study participants were seen for the randomization visit (month 0) and for scheduled visits at 2 weeks posttreatment initiation, monthly from months 1 to 4, and at months 6, 8, 10, and 12 for a total of 10 scheduled visits with monitoring for clinical malaria and toxicities. Caregivers were instructed to return to clinic or hospital whenever the child was unwell; all febrile children received malaria testing. After completing the blinded treatment phase, participants could receive open-label hydroxyurea, as per local Ethics Committee recommendations.

Measurements

Complete blood counts with leukocyte differential and absolute reticulocyte count (ARC) were measured at each visit (Sysmex Corporation, Kobe, Japan) plus blood chemistries using a Cobas 6000 analyzer Model C501 (Roche Diagnostics, Indianapolis, IN). HbF was quantified by capillary electrophoresis (Minicap; Sebia, Paris, France). Malaria was assessed by peripheral blood smear using Giemsa staining; each slide was read by 2 certified microscopists, with a third reading to resolve any discrepancies.

Clinical definitions

All children with measured fever (axillary temperature $\geq 37.5^{\circ}\text{C}$) in the clinic or fever by history were tested for malaria by microscopy. Children with measured fever or a history of fever and detectable *Plasmodium* species infection of any density on blood smear were diagnosed with clinical malaria, as is standard in this area. Malaria was treated with parenteral artesunate followed by oral artemether-lumefantrine, if the child was hospitalized, or oral artemether-lumefantrine, if the child was not hospitalized. The diagnosis of sickle-related clinical events followed published definitions,²⁴ with modifications such as pneumonia based on clinical rather than radiograph findings. Clinical sepsis was defined as presentation with fever and an unwell appearance, for which IV antibiotics were given. Full definitions for clinical adverse events (AEs) are provided in Table 1.

Study outcomes

The primary study outcome was incidence of clinical malaria. Secondary outcomes included the following: (1) a composite of 1 or more SCA-related AEs (pain, dactylitis, acute chest syndrome, splenic sequestration, or transfusion); (2) clinical AEs; and (3) dose-limiting toxicities.

Statistical analysis

We estimated malaria incidence would be between 0.3 and 1.3 malaria episodes per year, based on a study based in Kampala that includes community children

Table 2. Baseline demographic, clinical, and laboratory characteristics of the NOHARM randomized cohort

	Hydroxyurea (N = 104)	Placebo (N = 103)	P
Demographics, N (%)			
Age at enrollment, mean (SD), y	2.2 (0.9)	2.3 (0.9)	.3148
Male	55 (53)	57 (55)	.8298
Parent completing secondary school	32 (31)	35 (34)	.7614
Running water in the home	14 (13)	11 (11)	.6885
Growth measures, mean (SD)			
Height, cm	85.6 (8.8)	86.3 (8.2)	.5615
Weight, kg	11.3 (2.1)	11.5 (2.1)	.4604
Z score (weight for length/height)	−0.40 (1.06)	−0.39 (1.06)	.9316
Past medical history, N (%)			
Dactylitis	80 (77)	82 (80)	.7639
Vaso-occlusive crisis	88/103 (85)	81/103 (79)	.3421
Stroke	0	0	—
Splenomegaly	6/93 (6)	5/99 (5)	.7619*
Acute chest syndrome	21/103 (20)	13/100 (13)	.2219
Transfusion	56 (54)	57 (55)	.9393
Hospitalization within 1 y of enrollment	65 (63)	54 (52)	.1851
Laboratory measures, mean (SD)			
Hemoglobin, g/dL	7.5 (1.1)	7.6 (1.0)	.5214
MCV, fL	79 (9)	80 (9)	.8248
Fetal hemoglobin [HbF/(HbF + HbS)], %	14.6 (7.1)	13.3 (6.0)	.1591
ARC ($\times 10^9/L$)	380 (122)	381 (112)	.9623
WBC count, $\times 10^9/L$	19.0 (7.2)	18.7 (5.4)	.7148
ANC, $\times 10^9/L$	6.5 (3.1)	6.2 (2.6)	.4023
Platelets, $\times 10^9/L$	358 (171)	416 (138)	.0075
ALT, U/L	18 (9)	19 (9)	.5478
Creatinine, mg/dL	0.28 (0.09)	0.28 (0.07)	.6068

A total of 208 children were randomized to either hydroxyurea (N = 104) or placebo (N = 104). One participant was later deemed ineligible, and those data were removed from the data set. Values are shown as the mean (SD) or the number of study participants with the measure/total tested for that measure (% affected).

HbS, sickle hemoglobin; MCV, mean corpuscular volume; WBC, white blood cell.

*Fisher's exact test because of the low frequency.

(lowest incidence estimate) and children with severe malaria (highest incidence estimate) (C.C.J., unpublished data). With these rates of malaria incidence, and variance to mean ratio 1.7, we estimated that 100 children per treatment arm would provide 90% power ($\alpha = 0.05$) to detect between 59% and 34% difference in incidence between treatment arms, respectively. Study data were entered into OnCore and analyzed using R (Vienna, Austria; version 3.2.4). The primary analysis included all randomized participants per intention to treat, with $P < .05$ considered significant. Interim analysis for safety was performed after 100 children completed blinded study treatment. The difference in malaria incidence rate between the hydroxyurea and placebo groups was analyzed using negative binomial regression. For secondary outcomes, $P < .01$ was considered significant. Continuous measures are described as mean (standard deviation [SD]), and treatment arms were compared using the 2-sample Welch's t test, or when within the same individuals using a paired Welch's t test. Categorical measures are described using percent or frequency and compared using the Pearson χ^2 test with Yates correction or Fisher's exact test. Malaria episodes are reported as mean per child (95% confidence interval [CI]). Time to malaria infection is described by Kaplan-Meier curves. Groups were compared for cumulative incidence (ie, time to infection) using Gray's method, with all-cause death treated as a competing event.²⁵

Results


A total of 213 children were enrolled between 24 September 2014 and 2 October 2015; among these, 208 were randomized to either hydroxyurea or placebo (Figure 1). One child was later deemed ineligible because of an elevated baseline ALT, so that child's data were removed from the analyzed data set. The treatment arms had similar

baseline demographic, laboratory, and clinical measures (Table 2). No child had symptomatic or asymptomatic *Plasmodium* species parasitemia at enrollment.

Malaria occurred at a low rate throughout the study, perhaps reflecting excellent adherence to malaria prophylaxis. Over the 1-year study period, malaria testing was done for 235 episodes of measured fever or history of fever in study participants, with only 12 fever episodes because of malaria. Malaria incidence did not differ between children prescribed hydroxyurea (0.05 episodes per child per year, 95% CI [0.02, 0.13]) vs placebo (0.07 episodes per child per year [0.03, 0.16]); the hydroxyurea/placebo malaria incidence rate ratio was 0.7 ([0.2, 2.7], $P = .61$). Time to infection also did not differ significantly between hydroxyurea and placebo (Figure 2). All 12 malaria episodes were because of *P. falciparum*; 1 child had coinfection with *P. malariae*. Three children on hydroxyurea had a total of 5 malaria episodes, compared with 7 children on placebo with a total of 7 malaria episodes (Table 3). The median parasite density was 26 410 parasites per μL (minimum, 1064; maximum, 193 705). Six of the 12 episodes of malaria were severe, requiring hospitalization (2 with hemoglobin concentration < 5 g/dL, 1 with impaired consciousness, 1 with hemoglobin concentration < 5 g/dL and impaired consciousness, and 2 unable to take oral medication). All 10 study participants with malaria recovered. In the 12 confirmed cases of malaria, there were 4 episodes with concomitant clinical AE: vaso-occlusive pain crisis (N = 3) and acute chest syndrome/pneumonia (N = 1). In addition, there were 2 episodes of malaria with concomitant SAE: splenic sequestration (N = 1) and bacteremia (N = 1).

A per protocol composite clinical outcome included 1 or more SCA-related clinical events (vaso-occlusive painful crisis, dactylitis,

Figure 2. Incidence of malaria events in the NOHARM trial, with no statistical difference observed between the blinded treatment arms ($P = .19$). Comparisons of the hydroxyurea to placebo group were calculated using Gray's test for competing events, treating death as a competing event. Solid line represents hydroxyurea; dashed line represents placebo. The smaller inset diagram is identical to the larger graph but has a different scale.



acute chest syndrome/pneumonia, splenic sequestration, or blood transfusion). The proportion of children with this outcome was significantly lower in the hydroxyurea arm than the placebo arm (45% vs 69%; Table 3, $P = .001$). For individual clinical events, vaso-occlusive pain and hospitalizations were significantly less frequent with hydroxyurea than placebo (Table 3; Figure 3). This difference in hospitalization rate was largely driven by differences in vaso-occlusive crises, which accounted for 42% of all hospitalizations. The number needed to treat to prevent 1 hospitalization was 6.4, whereas the number needed to treat to prevent a SCA-related event was 2.5. No strokes occurred in either treatment arm during the 1-year treatment period. Clinical sepsis was more frequent with placebo (12.6%) than hydroxyurea (5.8%), but this difference was not statistically significant ($P = .14$). Blood cultures were positive in only 2 episodes, both for *Staphylococcus aureus*, 1 in each treatment arm.

SAEs, defined as death, an acute life-threatening event, or hospitalization for >7 days, occurred equally with 6 in each treatment arm (Table 3). During the blinded treatment phase, 3 children died (2 hydroxyurea, 1 placebo) and 5 others withdrew from the study. Causes of death were presumed sepsis (1 hydroxyurea, 1 placebo) and sudden death, cause unknown (hydroxyurea). In 2 of the SAEs listed in Table 3, 2 participants had malaria concomitant with bacteremia or splenic sequestration, as noted previously.

The 2 treatment arms had similar counts of laboratory AEs except anemia (Table 3). Specifically, low hemoglobin (<6.0 g/dL) occurred more frequently in children receiving placebo than hydroxyurea, whereas the frequencies of neutropenia, thrombocytopenia, and reticulocytopenia did not differ significantly between treatment arms. Similarly, the frequency of protocol-defined dose-limiting hematological toxicities did not differ between treatment arms (Table 3). Neutropenia was notably rare, occurring only once in only 2 participants over the entire study.

Children receiving hydroxyurea had significant treatment-associated increases in hemoglobin concentration, MCV, and HbF compared with those receiving placebo (Table 4). Substantial increases in HbF were observed for both children below and above the median enrollment age, despite starting with different baseline HbF levels. Conversely, children on hydroxyurea had significant decreases in WBC count, ANC, ARC, and platelets compared with children receiving placebo (Table 4). The treatment arms did not differ in changes in ALT or creatinine. Medication adherence, assessed by careful questioning at each clinic visit, was deemed excellent in both treatment arms.

Discussion

In this prospective randomized double-blinded placebo-controlled trial of young children with SCA living in Uganda, hydroxyurea therapy was both safe and efficacious. The 2 treatment arms did not differ in the incidence, severity, or other outcomes from malaria infection; in the incidence of clinical sepsis or bacteremia; or in the number of laboratory AEs or dose-limiting toxicities, including neutropenia. In contrast, the previously described laboratory and clinical benefits of hydroxyurea therapy were clearly observed in this young population. The increases in hemoglobin concentration and HbF levels, along with decreases in neutrophils and reticulocytes, were similar to values observed in US-based trials,^{4-7,26} and the rates of important clinical SCA-related events such as vaso-occlusive painful crisis and hospitalizations were similarly reduced. Taken together, these data suggest that hydroxyurea should be strongly considered as an important therapeutic option for young children with SCA living in malaria endemic areas.

NOHARM represents the first randomized trial of hydroxyurea in Africa. The study design captured all possible malaria events, with

Table 3. AEs in the NOHARM study population

	Hydroxyurea (N = 104)		Placebo (N = 103)		P
	Events	Participants	Events	Participants	
SAEs	6	6	6	6	
Bacteremia/sepsis	2	2	2	2	1.0*
Acute chest syndrome/pneumonia	1	1	2	2	.62*
Vaso-occlusive crisis	0	0	1	1	.50*
Acute splenic sequestration	2	2	0	0	.50*
Anemia	0	0	1	1	.50*
Sudden death	1	1	0	0	1.0*
SCA-related events (composite)					
Vaso-occlusive pain crisis, dactylitis, acute chest syndrome, splenic sequestration, or blood transfusion		47		71	.001
Clinical AEs	232	76	308	88	
Vaso-occlusive pain crisis/dactylitis	58	38	106	59	.004
Acute chest syndrome/pneumonia	24	21	32	24	.71
Clinical sepsis	8	6	16	13	.14
Acute splenic sequestration	0	0	0	0	—
Upper respiratory tract infection	107	54	108	62	.29
Gastrointestinal related	15	13	15	12	1.0
Malaria	5	3	7	7	.21*
Other infections	8	7	14	14	.16
Others (eg, injury)	7	7	10	9	.78
Clinical interventions	34	14	53	32	
Transfusion	14	12	18	17	.41
Hospitalization	20	12	35	28	.007
Laboratory AEs	57	31	78	46	
Anemia	40	25	65	42	.015
Reticulocytopenia	3	3	6	5	.50*
Neutropenia	2	2	0	0	.50*
Thrombocytopenia	12	11	4	4	.11
Elevated AST/ALT	0	0	2	1	.50*
Elevated bilirubin	0	0	1	1	.50*
Dose-limiting toxicities	21	15	17	13	
Anemia	6	4	8	8	.36
Reticulocytopenia	1	1	5	5	.12*
Neutropenia	2	2	0	0	.50*
Thrombocytopenia	12	11	4	4	.11

All serious adverse events (SAEs), AEs, and laboratory AEs were compared between treatment arms. Proportions of affected participants were compared by χ^2 tests with Yates correction. Full definitions of AEs are provided in Table 1.

*Fisher's exact test instead of χ^2 test, because of the low expected frequency.

blood smear testing for all episodes of current and recent fever. The incidence of malaria in our cohort was low, suggesting effective protection by insecticide-treated bed nets and monthly oral malaria prophylaxis with sulfadoxine-pyrimethamine provided to all study participants, or possibly a decreased risk of clinical malaria in children with SCA. Some form of malaria prophylaxis for children with SCA is a standard recommendation in most African countries,²⁷ and insecticide-treated bed net use for children <5 years of age in malaria endemic areas of Africa has increased dramatically over the past decade, to an average of >60% coverage,²⁸ so the preventive measures used in this study are similar to those in many other malaria endemic areas where children with SCA live. However, adherence to these recommendations is highly variable, so our study findings may not apply to children with SCA not on malaria prophylaxis and/or not using insecticide-treated bed nets. Because malaria incidence was low in our study, the range of relative differences in malaria incidence between hydroxyurea and placebo is broad (95% CI for incidence rate ratio 0.2 to 2.7), but the absolute differences in episodes is small (95% CI, 2-13 episodes per 100 children per year with hydroxyurea vs 3-16 episodes per 100 children per year with placebo). Hydroxyurea did not worsen malaria severity, because hospitalizations for malaria did not differ

between treatment arms, and no child died of malaria. Because of the low malaria incidence, this study does not provide definitive guidance regarding the safety of hydroxyurea in all malaria endemic areas, but the lack of increased risks of malaria incidence, severity, and outcome is reassuring. The incidence of malaria in these children with SCA further suggests that absolute differences in malaria risk with hydroxyurea will be very small. However, risks of hydroxyurea may differ in areas of higher malaria transmission, and genetic or environmental factors in other areas could affect risk of hydroxyurea toxicity or malaria-hydroxyurea interactions. In light of the present study data showing efficacy of hydroxyurea against pain crises and hospitalizations, with no evidence of increased malaria or infection risk in this area of low malaria transmission, it may be difficult to justify additional placebo-controlled trials in areas of higher malaria transmission. If additional placebo-controlled studies are not conducted, future studies of open-label hydroxyurea in other malaria endemic regions should carefully document the rates and complications of malaria during hydroxyurea treatment. These data will help determine the long-term safety profile of hydroxyurea for children living in malaria-endemic regions.

All 5 of the study children hospitalized with severe malaria (1 child was hospitalized twice) survived, in contrast to an earlier study from




Figure 3. Cumulative incidence of sickle-related AEs over 12 months, by blinded treatment arm. (A) Painful vaso-occlusive crisis; $P = .004$. (B) Hospitalization; $P = .002$. (C) Transfusion; $P = .27$. (D) Pneumonia/acute chest syndrome; $P = .51$.

Kenya in which 4 of 5 children with SCA hospitalized for severe malaria died (80% mortality).¹⁶ The presence of malaria parasites on peripheral blood smear also did not increase the risk of death during hospitalization, contrasting with a Tanzanian study of children with SCA, which reported increased mortality in hospitalized children with parasitemia.²⁹ In contrast, another study from Kenya reported no deaths among 38 children with SCA and severe malarial anemia, the most common form of severe malaria affecting these children.³⁰ Similarly, in a recent study in Kampala conducted by our group, none of 22 children with SCA who developed severe malarial anemia died.³¹ Together, these study findings cast doubt on the contention that severe malaria causes high rates of mortality in children with SCA. In the first 2 studies cited, which described increased mortality from malaria in children with SCA, malaria prophylaxis was either not given routinely¹⁶ or consisted of chloroquine.²⁹ The sulfadoxine-pyrimethamine prophylaxis given in the present study is likely more effective than chloroquine

prophylaxis. In addition, the 2 studies were conducted during a period of much lower bed net coverage, and a significant proportion of children in these earlier study cohorts received chloroquine or sulfadoxine-pyrimethamine for treatment of uncomplicated malaria, instead of the more effective current standard of care, artemisinin-combination therapy. Together, these factors may have played a role in the higher mortality seen with malaria in these prior cohorts.

Increased infection in children with drug-induced neutropenia was an additional concern regarding hydroxyurea treatment in an area where invasive bacterial infection is common in children. However, neutropenia was rare in NOHARM and did not differ on hydroxyurea vs placebo treatment, and episodes of pneumonia, clinically defined sepsis, and bacteremia did not differ between hydroxyurea or placebo treatment arms. These data confirm findings observed in the United States⁵ and should help allay safety concerns about hydroxyurea and infection risk related to neutropenia in low-resource settings.

Table 4. Selected laboratory characteristics of the NOHARM randomized population

Laboratory measures	Month 12			Change from baseline		
	Hydroxyurea	Placebo	P	Hydroxyurea	Placebo	P
Hemoglobin, g/dL	8.7 (1.3)	7.4 (1.0)	<.001	1.2 (1.2)	-0.1 (0.9)	<.001
MCV, fL	88 (9)	81 (8)	<.001	9 (7)	1 (5)	<.001
Fetal hemoglobin [HbF/(HbF + HbS)], %	22.9 (8.6)	10.4 (4.8)	<.001	8.5 (6.7)	-3.1 (3.3)	<.001
Enrollment age below median	24.1 (8.5)	12.1 (4.8)	<.001	7.9 (7.4)	-4.1 (3.5)	<.001
Enrollment age above median	21.4 (8.7)	8.8 (4.2)	<.001	9.3 (5.6)	-2.2 (2.8)	<.001
ARC, ×10 ⁹ /L	247 (107)	391 (122)	<.001	-144 (119)	11 (119)	<.001
WBC count, ×10 ⁹ /L	13.7 (5.1)	18.0 (5.1)	<.001	-5.4 (5.8)	-0.7 (5.1)	<.001
ANC, ×10 ⁹ /L	5.2 (2.5)	6.6 (2.6)	<.001	-1.3 (3.0)	0.4 (3.0)	<.001
Platelets, ×10 ⁹ /L	371 (166)	446 (143)	<.001	12 (184)	28 (152)	.50
ALT, U/L	19 (8)	18 (8)	.59	0.8 (11)	-0.9 (10)	.28
Creatinine, mg/dL	0.32 (0.11)	0.31 (0.08)	.64	0.04 (0.13)	0.04 (0.10)	.92

Results are shown only for participants who had data for the measure both at baseline and at 12 mo, as mean (SD) or the percentage of participants affected. The median age at enrollment was 2.2 y. Treatment group differences at month 12 and changes from baseline were tested by Welch's *t* tests.

Hydroxyurea was also associated with significantly fewer SCA-related clinical events, specifically vaso-occlusive crises, dactylitis, and hospitalizations. Other laboratory outcomes of long-term clinical importance for children with SCA, including increases in hemoglobin concentration and HbF, as well as decreases in leukocyte, neutrophil, and reticulocyte counts, were all more favorable in children receiving hydroxyurea than placebo, further supporting the drug's efficacy in this study population. Despite the differences in nutritional status from US-based populations, as evidenced by the low baseline *Z* scores, the hydroxyurea responses were remarkably similar to published results in the BABY HUG trial for both clinical and laboratory effects.⁵ The potential benefits of hydroxyurea treatment on growth in NOHARM are also important to assess but have not yet undergone formal analysis.

After the double-blinded treatment phase, the entire NOHARM cohort was offered open-label hydroxyurea as per the study design, and all but 1 family opted for treatment. Future analysis can thus compare the effects of early initiation and the longer-term effects of treatment. Because hydroxyurea escalated to maximum tolerated dose is standard in the United States,^{7,25} additional studies should assess how SCA-related AEs and hematological responses compare for children receiving fixed-dose treatment vs dose escalation regimens. Outcomes were excellent in the present study with a safe and relatively easily administered fixed dose (20 mg/kg), so it will be important to investigate how treatment benefits and risks differ with different dosing schemes of hydroxyurea for children with SCA living in malaria-endemic areas. Children in African countries often have limited access to health care facilities, and those facilities typically have limited laboratory testing capability. Less frequent laboratory monitoring would allow wider implementation of hydroxyurea treatment.

Additional studies that include longer duration of hydroxyurea in malaria-endemic areas will help determine optimal drug dosing, the range of AEs, and malaria incidence in areas with higher malaria transmission. Until then, the present study documents that with adequate monitoring, mosquito nets, and malaria prophylaxis, fixed-dose hydroxyurea treatment of children with SCA in malaria-endemic areas is safe with no increased incidence of malaria, sepsis, bacteremia, or SAEs. In combination with the observed clinical efficacy (fewer painful events and hospitalizations), the NOHARM study findings support the wider use of hydroxyurea for children with SCA living in malaria-endemic regions across sub-Saharan Africa.

Acknowledgments

The authors wish to thank the Mulago Hospital Sickle Cell Clinic and Global Health Uganda staff for conducting study work, the Data Coordinating Center staff at Cincinnati Children's Hospital Medical Center for setup of the study database and monitoring of study data, Fogarty and Doris Duke Charitable Foundation trainees who did operational work on the study, members of the Data and Safety Monitoring Board, and especially the study participants and their caregivers. Addmedica Inc. (Paris, France) donated both hydroxyurea and placebo study treatments.

This work was supported by a grant from the Doris Duke Charitable Foundation (ICRA 2013139), and Cincinnati Children's Research Foundation provided financial support for the NOHARM Data Coordinating Center.

The Doris Duke Charitable Foundation and Addmedica Inc. did not participate in the following aspects of the research: (1) study design; (2) data collection, analysis, or interpretation; (3) writing the report; or (4) decision to submit the publication.

Authorship

Contribution: R.O.O., R.E.W., and C.C.J. designed the study, supervised the trial, analyzed the results, and wrote the first draft of the manuscript; T.S.L. helped coordinate many critical aspects of the trial to ensure its safe and successful operational execution; C.M.N., H.A.H., and P.K. enrolled patients, collected data, and helped interpret the results; A.L. and J.S.H. performed statistical analyses for the trial; and all authors participated in the editing of the manuscript and approved the final version.

Conflict-of-interest disclosure: R.E.W. is a consultant for Global Blood Therapeutics and Nova Laboratories, is on an advisory board for Agios Pharmaceuticals, receives research support from Bristol-Myers Squibb, and serves on a Data and Safety Monitoring Board for the US Food and Drug Administration. None of these disclosures are relevant to the results and conclusions of the NOHARM trial. The remaining authors declare no competing financial interests.

ORCID profiles: R.E.W., 0000-0001-9582-0594; C.C.J., 0000-0002-1634-1411.

Correspondence: Russell E. Ware, Division of Hematology, Department of Pediatrics, Cincinnati Children's Hospital Medical Center, 3333 Burnet Ave, Cincinnati, OH 45229; e-mail: russell.ware@cchmc.org.

References

- Piel FB, Patil AP, Howes RE, et al. Global distribution of the sickle cell gene and geographical confirmation of the malaria hypothesis. *Nat Commun*. 2010;1(8):104-111.
- Grosse SD, Odame I, Atrash HK, Amendah DD, Piel FB, Williams TN. Sickle cell disease in Africa: a neglected cause of early childhood mortality. *Am J Prev Med*. 2011;41(6, suppl 4):S398-S405.
- Allison AC. Protection afforded by sickle-cell trait against subtertian malarial infection. *BMJ*. 1954;1(4857):290-294.
- Charache S, Terrin ML, Moore RD, et al; Investigators of the Multicenter Study of Hydroxyurea in Sickle Cell Anemia. Effect of hydroxyurea on the frequency of painful crises in sickle cell anemia. *N Engl J Med*. 1995;332(20):1317-1322.
- Kinney TR, Helms RW, O'Branski EE, et al; Pediatric Hydroxyurea Group. Safety of hydroxyurea in children with sickle cell anemia: results of the HUG-KIDS study, a phase I/II trial. *Blood*. 1999;94(5):1550-1554.
- Wang WC, Ware RE, Miller ST, et al; BABY HUG investigators. Hydroxycarbamide in very young children with sickle-cell anaemia: a multicentre, randomised, controlled trial (BABY HUG). *Lancet*. 2011;377(9778):1663-1672.
- Ware RE, Davis BR, Schultz WH, et al. Hydroxycarbamide versus chronic transfusion for maintenance of transcranial doppler flow velocities in children with sickle cell anaemia-TCD with transfusions changing to hydroxyurea (TWITCH): a multicentre, open-label, phase 3, non-inferiority trial. *Lancet*. 2016;387(10019):661-670.
- Ware RE. How I use hydroxyurea to treat young patients with sickle cell anemia. *Blood*. 2010;115(26):5300-5311.
- Yawn BP, Buchanan GR, Afenyi-Annan AN, et al. Management of sickle cell disease: summary of the 2014 evidence-based report by expert panel members. *JAMA*. 2014;312(10):1033-1048.
- Brun M, Bourdoulous S, Couraud PO, Elion J, Krishnamoorthy R, Lapoumeroulie C. Hydroxyurea downregulates endothelin-1 gene expression and upregulates ICAM-1 gene expression in cultured human endothelial cells. *Pharmacogenomics J*. 2003;3(4):215-226.
- Berendt AR, Simmons DL, Tansey J, Newbold CI, Marsh K. Intercellular adhesion molecule-1 is an endothelial cell adhesion receptor for *Plasmodium falciparum*. *Nature*. 1989;341(6237):57-59.
- Navarra P, Grohmann U, Nocentini G, et al. Hydroxyurea induces the gene expression and synthesis of proinflammatory cytokines in vivo. *J Pharmacol Exp Ther*. 1997;280(1):477-482.
- Grau GE, Taylor TE, Molyneux ME, et al. Tumor necrosis factor and disease severity in children with falciparum malaria. *N Engl J Med*. 1989;320(24):1586-1591.
- Conran N, Fattori A, Saad ST, Costa FF. Increased levels of soluble ICAM-1 in the plasma of sickle cell patients are reversed by hydroxyurea. *Am J Hematol*. 2004;76(4):343-347.
- Pino P, Taoufiq Z, Brun M, et al. Effects of hydroxyurea on malaria, parasite growth and adhesion in experimental models. *Parasite Immunol*. 2006;28(12):675-680.
- McAuley CF, Webb C, Makani J, et al. High mortality from *Plasmodium falciparum* malaria in children living with sickle cell anemia on the coast of Kenya. *Blood*. 2010;116(10):1663-1668.
- Ramakrishnan M, Moisi JC, Klugman KP, et al. Increased risk of invasive bacterial infections in African people with sickle-cell disease: a systematic review and meta-analysis. *Lancet Infect Dis*. 2010;10(5):329-337.
- Amaratunga C, Lopera-Mesa TM, Brittain NJ, et al. A role for fetal hemoglobin and maternal immune IgG in infant resistance to *Plasmodium falciparum* malaria. *PLoS One*. 2011;6(4):e14798.
- Cokic VP, Smith RD, Beleslin-Cokic BB, et al. Hydroxyurea induces fetal hemoglobin by the nitric oxide-dependent activation of soluble guanylyl cyclase. *J Clin Invest*. 2003;111(2):231-239.
- Cabrales P, Zanini GM, Meays D, Frangos JA, Carvalho LJ. Nitric oxide protection against murine cerebral malaria is associated with improved cerebral microcirculatory physiology. *J Infect Dis*. 2011;203(10):1454-1463.
- Serghides L, Kim H, Lu Z, et al. Inhaled nitric oxide reduces endothelial activation and parasite accumulation in the brain, and enhances survival in experimental cerebral malaria. *PLoS One*. 2011;6(11):e27714.
- Anstey NM, Weinberg JB, Hassanali MY, et al. Nitric oxide in Tanzanian children with malaria: inverse relationship between malaria severity and nitric oxide production/nitric oxide synthase type 2 expression. *J Exp Med*. 1996;184(2):557-567.
- Anyanwu JN, Williams O, Sautter CL, et al. Novel use of hydroxyurea in an African region with malaria: protocol for a randomized controlled clinical trial. *JMIR Res Protoc*. 2016;5(2):e110.
- Ballas SK, Lief S, Benjamin LJ, et al; Investigators, Comprehensive Sickle Cell Centers. Definitions of the phenotypic manifestations of sickle cell disease. *Am J Hematol*. 2010;85(1):6-13.
- Gray RJ. A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Ann Stat*. 1988;16(3):1141-1154.
- Ware RE. Optimizing hydroxyurea therapy for sickle cell anemia. *Hematology Am Soc Hematol Educ Program*. 2015;2015(1):436-443.
- Aneni EC, Hamer DH, Gill CJ. Systematic review of current and emerging strategies for reducing morbidity from malaria in sickle cell disease. *Trop Med Int Health*. 2013;18(3):313-327.
- UNICEF. Malaria mortality among children under five is concentrated in sub-Saharan Africa. <http://data.unicef.org/topic/child-health/malaria/#>. Accessed 26 July 2017.
- Makani J, Komba AN, Cox SE, et al. Malaria in patients with sickle cell anemia: burden, risk factors, and outcome at the outpatient clinic and during hospitalization. *Blood*. 2010;115(2):215-220.
- Komba AN, Makani J, Sadarangani M, et al. Malaria as a cause of morbidity and mortality in children with homozygous sickle cell disease on the coast of Kenya. *Clin Infect Dis*. 2009;49(2):216-222.
- Opoka RO, Bangirana P, Idro R, Shabani E, Namazzi R, John CC. Lack of mortality in 22 children with sickle cell anemia and severe malarial anemia [published online ahead of print 17 August 2017]. *Pediatr Blood Cancer*. doi:10.1002/pbc.26745.



blood[®]

2017 130: 2585-2593

doi:10.1182/blood-2017-06-788935 originally published
online October 19, 2017

Novel use Of Hydroxyurea in an African Region with Malaria (NOHARM): a trial for children with sickle cell anemia

Robert O. Opoka, Christopher M. Ndugwa, Teresa S. Latham, Adam Lane, Heather A. Hume, Phillip Kasirye, James S. Hodges, Russell E. Ware and Chandy C. John

Updated information and services can be found at:

<http://www.bloodjournal.org/content/130/24/2585.full.html>

Articles on similar topics can be found in the following Blood collections

[Clinical Trials and Observations](#) (4681 articles)

[Free Research Articles](#) (4827 articles)

[Pediatric Hematology](#) (546 articles)

[Plenary Papers](#) (507 articles)

[Red Cells, Iron, and Erythropoiesis](#) (845 articles)

[Sickle Cell Disease](#) (140 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>